

FREE GUIDE

Postpartum Recovery

Physical and emotional recovery, week by week — and when to call your doctor.

By Anahely

BYANAHEL.Y.COM

A NOTE FROM ANAHELY

The pamphlet nobody actually gives you.

Nobody really prepared me for the fourth trimester. Pregnancy gets all the books; recovery gets a pamphlet on your way out the door. This is everything I wish that pamphlet actually said.

NOT A SUBSTITUTE FOR MEDICAL CARE

This guide is general guidance, not medical advice. If something feels wrong, trust that instinct and call your provider — you don't need to justify the call.

Physical Recovery

Recovery isn't linear, and it isn't the same for a vaginal birth versus a C-section — but here's a rough shape of what the weeks tend to look like.

WEEK 1–2

Bleeding (lochia) is heaviest, similar to a heavy period, and gradually tapers. Cramping as your uterus contracts back down. Perineal soreness or incision pain, depending on delivery. Rest is not optional here — your body is doing enormous repair work.

WEEK 3–6

Bleeding lightens to spotting, then usually stops. Energy starts returning in small windows. Most providers clear light activity around 6 weeks, but “cleared” doesn't mean “back to normal” — many people need longer.

WEEK 6–12

Hormones are still shifting. Hair shedding often starts around month 3–4 (temporary). Core and pelvic floor strength continues rebuilding — this is a good window to ask about pelvic floor physical therapy if anything feels off.

A padsicle (a maxi pad chilled with witch hazel and aloe) sounds like a Pinterest gimmick, but it genuinely helped every single time.

Emotional Recovery

The “baby blues” are extremely common and different from postpartum depression — knowing the difference matters.

Usually the Baby Blues

- Crying for no clear reason
- Feeling overwhelmed or irritable
- Mood swings that come and go
- Typically peaks day 3-5, resolves within 2 weeks

Worth Telling Your Provider

- Sadness lasting more than 2 weeks
- Difficulty bonding with the baby
- Intrusive or scary thoughts
- Thoughts of harming yourself

Postpartum depression and anxiety are medical conditions, not a character flaw — they're also very treatable. Say the hard sentence out loud to your provider.

What Actually Helped

Small, unglamorous things that made a real difference across five recoveries.

A “recovery station”

Water bottle, snacks, charger, and pads all within arm's reach of wherever you'll be feeding baby most.

Saying yes to help

Let someone else hold the baby so you can shower. It felt indulgent every time; it never was.

A peri bottle, always

Use it every single bathroom trip for the first couple weeks — genuinely one of recovery's best-kept secrets.

Lowering the bar

The goal most days was “everyone fed, everyone safe.” Some days that was the whole win, and that was enough.

When to Call Your Doctor

Postpartum complications are uncommon but serious — these are always worth a call, not a wait-and-see.

NOT A SUBSTITUTE FOR MEDICAL CARE

If something feels wrong, trust that instinct and call — you don't need to justify the call.

Usually Normal

- Bleeding that gradually lightens over 4–6 weeks
- Mild cramping, especially while nursing
- Some swelling in feet/hands the first week
- Emotional ups and downs in first two weeks

Call Your Doctor

- Bleeding that soaks a pad in an hour, or clots
- Fever over 100.4°F (38°C)
- Redness/discharge at an incision site
- Chest pain, shortness of breath, leg swelling
- Thoughts of harming yourself or the baby

*"Recovery isn't a race back to who you were —
it's the slow work of becoming who you are now."*

MORE GUIDES AT [BYANAHEL.Y.COM](https://byanahely.com)

@byanahely on Instagram